COMMENTARY

Common but differentiated **IBI Prize Essav**

1056



LETTERS I BOOKS I POLICY FORUM I EDUCATION FORUM I PERSPECTIVES

LETTERS

edited by Jennifer Sills

The Scientific Whaling Loophole

AT THE RECENT INTERNATIONAL WHALING COMMISSION'S ANNUAL meeting in Panama, South Korean officials announced their plan to initiate a "scientific whaling" program (1). This announcement came as a surprise given the general sentiment that the global demand for whale meat is declining. After weeks of international outcry, on 17 July, South Korea reversed their decision to hunt whales for research, but the issue is not dead (2).

South Korea claimed that the goal of the scientific whaling pro-

gram is to study the types and amounts of fish whales eat, given conflict with fisheries. Yet, it is well established in the scientific literature that there are many ways to study whale diet without killing them (3).

Decades of fruitless negotiation between pro- and anti-whaling nations suggests a broken system, wrought with loopholes that allow unsustainable whaling to continue. Within this broken system, there is no incentive to reduce whaling, as the recent announcement by South Korea shows. Whaling groups are unwilling to compromise by allowing a sustainable harvest of whales, so unsustainable (scientific) whaling continues.

To ensure a future of both whales and whalers, we must harness the passion and value that people place on living whales, without telling people what to do or force one set of values on others. We need novel, out-of-the-box approaches to effective management and conservation of whales. We must compromise to ensure reductions in whales being killed, better oversight of countries that harvest them, and limited whaling that does not threaten the persistence of whales.

For those who believe that whaling is unethical, I challenge you to put forward alternative ideas to a global moratorium that fosters the "loophole" of scientific whaling. With new plans to develop scientific whaling programs (4), the current global moratorium is clearly broken. Scientists, conservation advocates, resource managers, and the public must work together to develop new approaches to ensure the persistence of whales in our oceans.

LEAH R. GERBER

Faculty of Ecology, Evolution, and Environmental Sciences, School of Life Sciences, Arizona State University, Tempe, AZ 85281, USA. E-mail: leah.gerber@asu.edu

References

- 1. Associated Press, The New York Times, 6 July 2012, p. A7 (www.nytimes.com/2012/07/06/ world/europe/south-korea-whaling-plan-criticized.html).
- 2. S. Herman, Voice of America, 24 July 2012 (www.voanews.com/content/south-korea-stillconsidering-resumption-of-whaling/1444081.html).
- 3. L. R. Gerber, L. Morissette, K. Kaschner, D. Pauly, Science 323, 880 (2009).
- 4. R. Black, BBC News, 5 July 2012 (www.bbc.co.uk/news/science-environment-18719512).

Iconic CO₂ Time Series at Risk

THE STEADY RISE IN ATMOSPHERIC LONGlived greenhouse gas concentrations is the main driver of contemporary climate change. The Mauna Loa CO₂ time series (1, 2), started by C. D. Keeling in 1958 and maintained today by the Scripps Institution of Oceanography and the Earth System Research Laboratory (ESRL) of NOAA, is iconic evidence of the effect of humancaused fossil fuel and land-use change emissions on the atmospheric increase of CO₂. The continuity of such records depends critically on having stable funding, which is challenging to maintain in the context of 3- to 4-year research grant funding cycles (3), and is currently threatened by the financial crisis.

The ESRL Global Monitoring Division maintains a network of about 100 surface and aircraft sites worldwide at which whole air samples are collected approximately every week for analysis of CO₂, CH₄, CO, halocarbons, and many other chemical species (4). This is complemented by high-frequency measurements at the Mauna Loa, Barrow, American Samoa, and South Pole observatories, and about 10 North American tall towers. The success of the NOAA program has inspired similar efforts in Europe (5), China (6), India (7), and Brazil (8), with the United Nations World Meteorological Organization providing guidance and precision requirements through the Global Atmosphere Watch program (9), but no funding.

The data collected by NOAA and its worldwide partners have been used not only to demonstrate the unassailable rise of atmospheric greenhouse gas concentrations, but also to infer the magnitudes, locations, and times of surface-atmosphere exchange of those gases based on small concentration gradients between sites (10). Important findings from analysis of these records include the detection of a significant terrestrial carbon sink at northern mid-latitudes (11) and subsequent research aimed at identifying the mechanisms by which that sink must operate. Long-term, high-quality, atmospheric measurements are crucial for quantifying trends in greenhouse gas fluxes and attributing them to fossil fuel emissions, changes in land-use and management, or the response of natural

DIT

land and ocean ecosystems to climate change and elevated CO_2 concentrations.

Greenhouse gas measurements along tall towers in the interior continents allow quantification of regional sources and sinks, which has a very high relevance for measuring the effectiveness of climate policy. NOAA ESRL provides measurements that are critical for the U.S. national security in that they provide independent verification and early warning of changing greenhouse gas emissions from countries involved in efforts to mitigate greenhouse gases.

Dedicated carbon-observing satellites such as GOSAT and OCO-2 are needed to fill in the missing geographical information required for verification of carbon flux mitigation efforts. However, satellite retrievals do not yet provide sufficient information to deliver new constraints on surface fluxes, although quick progress is being made in this direction. In situ observations are crucial for anchoring space-borne measurements, for detecting potential biases of remote sensing techniques, and for providing continuity given the finite lifetime of satellites.

Despite the growing importance of greenhouse gas observations to humanity, sub-

stantial budget cuts at NOAA have resulted in curtailment of our ability to observe and understand changes to the global carbon cycle. Already, a dozen surface flasksampling sites have been removed from NOAA's operational network and aircraft profiling sites have been eliminated and reduced in frequency at the remaining NOAA sites. The planned growth in the tall tower program has stopped, and plans for closing some towers are being developed. The U.S. budget process in this election year, with the added risk of mandatory across-the-board cuts due to the 2011 Budget Control Act, foretells more bleak news for greenhouse gas monitoring at NOAA and could cause further retreat from the goal of recording ongoing changes in atmospheric composition. As scientists, we believe that preserving the continuity of these vital time series must remain a priority for U.S. carbon cycle research.

SANDER HOUWELING,^{1,2*} BAKR BADAWY,³ DAVID F. BAKER,⁴ SOURISH BASU,^{1,2} DMITRY BELIKOV,⁵ PETER BERGAMASCHI,⁶ PHILIPPE BOUSQUET,⁷ GREGOIRE BROQUET,⁷ TIM BUTLER,⁸ JOSEP G. CANADELL,⁹ JING CHEN,¹⁰ FREDERIC CHEVALLIER,⁷ PHILIPPE CIAIS,⁷ G. JAMES COLLATZ,¹¹ SCOTT DENNING,⁴

RICHARD ENGELEN,¹² IAN G. ENTING,¹³ MARC L. FISCHER, 14 ANNEMARIE FRASER, 15 CHRISTOPH GERBIG,³ MANUEL GLOOR,¹⁶ ANDREW R. JACOBSON, 17,18 DYLAN B. A. JONES, 10 MARTIN HEIMANN,³ ASLAM KHALIL,¹⁹ THOMAS KAMINSKI,²⁰ PRASAD S. KASIBHATLA,²¹ NIR Y. KRAKAUER,²² MAARTEN KROL,^{1,2,23} TAKASHI MAKI,24 SHAMIL MAKSYUTOV,5 ANDREW MANNING,25 ANTOON MEESTERS,26 JOHN B. MILLER, 17,18 PAUL I. PALMER, 15 PRABIR PATRA,²⁷ WOUTER PETERS,²³ PHILIPPE PEYLIN,⁷ ZEGBEU POUSSI,²⁸ MICHAEL J. PRATHER, 29 JAMES T. RANDERSON, 29 THOMAS RÖCKMANN,² CHRISTIAN RÖDENBECK,³ JORGE L. SARMIENTO, 30 DAVID S. SCHIMEL, 31 MARKO SCHOLZE, 32 ANDREW SCHUH, 4 PARV SUNTHARALINGAM,²⁵ TARO TAKAHASHI,³³ JOCELYN TURNBULL,³⁴ LEONID YURGANOV,³⁵ ALEX VERMEULEN³⁶

¹SRON Netherlands Institute for Space Research, 3584 CA, Utrecht, Netherlands. ²Institute for Marine and Atmospheric Research Utrecht, 3584 CC Utrecht, Netherlands. ³Max-Planck-Institute for Biogeochemistry, 07745, Jena, Germany. ⁴Cooperative Institute for Research in the Atmosphere, Colorado State University, Fort Collins, CO 80523– 1375, USA. ⁵National Institute for Environmental Studies, Tsukuba, 305-8506, Japan. ⁴European Commission Joint Research Centre, Ispra, Italy. ⁷Laboratoire des Sciences du Climat et de l'Environnement, Unité mixte CEA, UVSQ, CNRS, 91191, Gif-sur-Yvette, France. ⁸Institute for Advanced Sustainability Studies, 14467, Potsdam, Germany.

AAAS Award for Science Diplomacy

Many scientists and engineers contribute valuable time away from the established career paths of research, teaching, and publishing to foster activities and develop programs that both address key science questions and build important societal links. AAAS seeks to recognize an individual or a limited number of individuals working together in the scientific or engineering community for making an outstanding contribution to furthering science diplomacy.

The recipient receives US \$5,000 award, a commemorative plaque, complimentary registration, and reimbursement for reasonable travel and hotel expenses.

The award is open to all regardless of nationality or citizenship. Nominees must be living at the time of their nomination. Please visit <u>http://www.aaas.org/aboutaaas/awards/int/</u> for more information and nomination instructions.

All materials must be received by September 1.





⁹Global Carbon Project, Commonwealth Scientific and Industrial Research Organisation (CSIRO), Canberra, ACT 2601, Australia. ¹⁰University of Toronto, Toronto, ON, M5S 1A7, Canada. ¹¹NASA's Goddard Space Flight Center, Greenbelt, MD 20771, USA. ¹²European Centre for Medium-Range Weather Forecasts, Shinfield Park, Reading, RG2 9AX, UK. ¹³ARC Centre of Excellence in the Mathematics and Statistics of Complex Systems, University of Melbourne, Victoria 3010, Australia. ¹⁴Lawrence Berkeley National Laboratory, Washington, DC 20024, USA. 15University of Edinburgh, Edinburgh, EH9 3JW, UK. ¹⁶Earth and Biosphere Institute, School of Geography, University of Leeds, Leeds LS2 9]T, UK. 17Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, CO 80309, USA. 18NOAA Earth System Research Laboratory, Boulder, CO 80305, USA. 19Portland State University, Portland, OR 97207, USA. 20FastOpt, 22767 Hamburg, Germany. 21Nicholas School of the Environment, Duke University, Durham NC 27708, USA. ²²Department of Civil Engineering, City College of New York, New York, NY 10031, USA. ²³Meteorology and Air Quality, Wageningen University and Research Center, 6708 PB Wageningen, Netherlands. ²⁴Environmental and Applied Meteorology Research Department, Meteorological Research Institute, Tskuba, Japan. 25 University of East Anglia, Norwich Research Park, Norwich, NR4 7TJ, UK. ²⁶Vrije Universiteit, 1081 HV, Amsterdam, Netherlands. ²⁷Research Institute for Global Change, Japan Agency for Marine-Earth Science and Technology, Yokohama, 236-0001, Japan. ²⁸ClimMod, 91401 Orsay, France. ²⁹University of California Irvine, Irvine, CA 92697, USA. ³⁰Princeton University, Princeton, NJ 08544, USA. ³¹National Ecological Observatory Inc., Boulder, CO 80301, USA. 32 Bristol University, Clifton, BS8 1RJ, UK. ³³Lamont-Doherty Earth Observatory, Columbia University, Palisades, NY 10964-8000, USA. ³⁴National Isotope Centre, GNS Science, Lower Hutt, New Zealand. ³⁵University of Maryland Baltimore County, Baltimore, MD 21250, USA. ³⁶Energieonderzoek Centrum Nederland, 1755 ZG Petten, Netherlands.

*To whom correspondence should be addressed. E-mail: s.houweling@sron.nl

References and Notes

- 1. Scripps CO₂ Program (http://scrippsco2.ucsd.edu).
- ESRL Global Monitoring Division, Recent Mauna Loa CO₂ (www.esrl.noaa.gov/gmd/ccgg/trends).
- 3. C. D. Keeling, Annu. Rev. Energy Environ. 23, 25 (1998).
- 4. ESRL Global Monitoring Division, GMD Carbon Cycle
- Greenhouse Gases Group (www.esrl.noaa.gov/gmd/ccgg). 5. Integrated Carbon Observing System (www.icos-
- infrastructure.eu).6. China Meteorological Association (www.cngaw-ghgs.org) [in Chinese].
- W. A. Brand, Ed., World Meteorological Organization, Global Atmosphere Watch, GAW Report No. 194, WMO/ TD-No. 1553 (2011); www.wmo.int/pages/prog/arep/gaw/ documents/GAW_194_WMO_TD_1553_web.pdf.
- 8. L. Gatti et al., Tellus Ser. B 62, 581 (2010).
- WMO Global Atmosphere Watch, World Data Centre for Greenhouse Gases (www.wmo.int/pages/prog/arep/gaw/ gaw_home_en.html).
- 10. K. R. Gurney et al., Nature 415, 626 (2002).
- 11. P. P. Tans et al., Science 247, 1431 (1990).
- 12. The authors collaborate in the international transport model intercomparison project, TRANSCOM.

Letters to the Editor

Letters (~300 words) discuss material published in *Science* in the past 3 months or matters of general interest. Letters are not acknowledged upon receipt. Whether published in full or in part, Letters are subject to editing for clarity and space. Letters submitted, published, or posted elsewhere, in print or online, will be disqualified. To submit a Letter, go to www.submit2science.org.

Decoding Cryptosystems

R. STONE'S NEWS FOCUS STORY ABOUT PAN Jianwei's marvelous quantum optics experiments ("Entangled secret messages from space," 29 June, p. 1632) propagates some unfortunately common misconceptions about the uses of quantum key distribution (QKD) technology, especially its integration into a complete cryptosystem.

The confusion arises in the distinction between a cryptographic key and a communication session encrypted via the key. QKD does not carry or encrypt the message directly. Instead, QKD uses a classical communication authentication mechanism, quantum eavesdropping detection, and a healthy dose of statistics, as well as both quantum and classical randomness, to generate a random bit string that is known to be secret and shared only between two parties. This random bit string is then used as the encryption key for a standard, classical encryption system.

The ultimate success of the cryptosystem in protecting sensitive data depends on several factors. One such factor is the QKD implementation itself; no general attack against QKD is known, but various attacks have been proposed and even implemented against the photon generators, detectors, and electromechanical subsystems. Implementers respond by fixing problems in the usual thrust-andparry of security systems implementation.

The security of the data depends on the strength of the classical encryption. The ideal use of the key would be to use it once and discard, as in a one-time pad (OTP), but current QKD key generation rates are far below desired classical communication rates, leading implementers to use the key for encryption schemes, such as Advanced Encryption Standard (AES), which encrypt many data bits with the use of fewer key bits. If used properly, OTP is perfectly secure, whereas AES could be broken by trying all possible keys, a theoretically possible but computationally impractical task.

Rather than flat statements that communication using QKD is totally unbreakable, it is more correct to say that it presents a different attack surface.

RODNEY VAN METER

Faculty of Environment and Information Studies, Keio University, Fujisawa, Kanagawa 252-0882, Japan. E-mail: rdv@sfc.wide.ad.jp

CORRECTIONS AND CLARIFICATIONS

News Focus: "Where are the missing baryons?" by Y. Bhattacharjee (1 June, p. 1093). Oxygen VI is oxygen stripped of five electrons, not six, and Neon VIII is neon stripped of seven electrons, not eight. **This Week in Science:** "Tic TOC1 plant clock" (6 April, p. 11). The editors note that the title of this summary was not intended to convey a connection between TOC1 and the plant gene *Tic*.

Reports: "The B73 maize genome: Complexity, diversity, and dynamics" by P. S. Schnable *et al.* (20 November 2009, p. 1112). Reference 27 should be C. Liang, L. Mao, D. Ware, L. Stein, *Genome Res.* **19**, 1912 (2009). The reference has been corrected in the HTML version online.

TECHNICAL COMMENT ABSTRACTS

Comment on "Intensifying Weathering and Land Use in Iron Age Central Africa"

K. Neumann, M. K. H. Eggert, R. Oslisly, B. Clist, T. Denham, P. de Maret, S. Ozainne, E. Hildebrand, K. Bostoen, U. Salzmann, D. Schwartz, B. Eichhorn, B. Tchiengué, A. Höhn Bayon *et al.* (Reports, 9 March 2012, p. 1219) interpreted unusually high aluminum-potassium ratio values in an Atlantic sediment core as indicating anthropogenic deforestation around 2500 years before the present (B.P.). We argue that there is no terrestrial evidence for forest destruction by humans and that the third millennium B.P. rainforest crisis can be clearly attributed mostly to climatic change.

Full text at www.sciencemag.org/cgi/content/full/337/ 6098/1040-c

Comment on "Intensifying Weathering and Land Use in Iron Age Central Africa"

Jean Maley, Pierre Giresse, Charles Doumenge, Charly Favier

Bayon *et al.* (Reports, 9 March 2012, p. 1219) claim that the "rainforest crisis" in Central Africa centered around 2500 years before the present "was not triggered by natural climatic factors" and that it was caused by widespread deforestation resulting from the arrival of the Bantu colonists. However, there is a consensus among palaeoecologists that this landscape change and the related physical erosion it caused was due mainly to a shift to more seasonal rainfall regime.

Full text at www.sciencemag.org/cgi/content/full/337/ 6098/1040-d

Response to Comments on "Intensifying Weathering and Land Use in Iron Age Central Africa"

Germain Bayon, Bernard Dennielou, Joël Etoubleau, Emmanuel Ponzevera, Samuel Toucanne, Sylvain Bermell

Neumann *et al.* argue that terrestrial evidence does not support our interpretation of large-scale human land use in Central Africa about 2500 years ago and that climate was the main driver of the rainforest crisis at that time, and Maley *et al.* raise a number of concerns about our interpretation of data from chemical weathering proxies. Taking into account existing palaeoclimatic data and clarifying some misconceptions, we reassert that humans must also have contributed fundamentally to this large vegetation change.

Full text at www.sciencemag.org/cgi/content/full/337/ 6098/1040-e

31 AUGUST 2012 VOL 337 SCIENCE www.sciencemag.org





Iconic CO₂ Time Series at Risk Sander Houweling, Bakr Badawy, David F. Baker, Sourish Basu, Dmitry Belikov, Peter Bergamaschi, Philippe Bousquet, Gregoire Broquet, Tim Butler, Josep G. Canadell, Jing Chen, Frederic Chevallier, Philippe Ciais, G. James Collatz, Scott Denning, Richard Engelen, Ian G. Enting, Marc L. Fischer, Annemarie Fraser, Christoph Gerbig, Manuel Gloor, Andrew R. Jacobson, Dylan B. A. Jones, Martin Heimann, Aslam Khalil, Thomas Kaminski, Prasad S. Kasibhatla, Nir Y. Krakauer, Maarten Krol, Takashi Maki, Shamil Maksyutov, Andrew Manning, Antoon Meesters, John B. Miller, Paul I. Palmer, Prabir Patra, Wouter Peters, Philippe Peylin, Zegbeu Poussi, Michael J. Prather, James T. Randerson, Thomas Röckmann, Christian Rödenbeck, Jorge L. Sarmiento, David S. Schimel, Marko Scholze, Andrew Schuh, Parv Suntharalingam, Taro Takahashi, Jocelyn Turnbull, Leonid Yurganov and Alex Vermeulen (August 30. 2012)

Science 337 (6098), 1038-1040. [doi: 10.1126/science.337.6098.1038-b]

Editor's Summary

This copy is for your personal, non-commercial use only.

Article Tools	Visit the online version of this article to access the personalization and article tools: http://science.sciencemag.org/content/337/6098/1038.2
Permissions	Obtain information about reproducing this article: http://www.sciencemag.org/about/permissions.dtl

Science (print ISSN 0036-8075; online ISSN 1095-9203) is published weekly, except the last week in December, by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. Copyright 2016 by the American Association for the Advancement of Science; all rights reserved. The title *Science* is a registered trademark of AAAS.