



# Positive impacts of precipitation intensity on monthly CO<sub>2</sub> fluxes in North America

Chaoyang Wu<sup>a,b,\*</sup>, Jing M. Chen<sup>b</sup>, Ankur R. Desai<sup>c</sup>, Peter M. Lafleur<sup>d</sup>, Shashi B. Verma<sup>e</sup>

<sup>a</sup> The State Key Laboratory of Remote Sensing Science, Laboratory of Digital Earth Sciences, Institute of Remote Sensing and Digital Earth, Chinese Academy of Sciences, Beijing, China

<sup>b</sup> Department of Geography, University of Toronto, 100 St. George St., Room 5047, Toronto, ON, Canada M5S 3G3

<sup>c</sup> Dept. of Atmospheric and Oceanic Sciences, University of Wisconsin – Madison, 1225 W Dayton St., Madison, WI 53706 USA

<sup>d</sup> Department of Geography, Trent University, Peterborough, Ontario, Canada K9J 7B8

<sup>e</sup> School of Natural Resources, University of Nebraska-Lincoln, 807 Hardin Hall, 3310 Holdrege Street, Lincoln, NE, USA

## ARTICLE INFO

### Article history:

Received 27 November 2011

Accepted 25 October 2012

Available online 1 November 2012

### Keywords:

Precipitation pattern

Gross ecosystem production

Leaf area index

North America

## ABSTRACT

Precipitation is one of the most important climate factors that can affect the gross ecosystem production (GEP) of terrestrial ecosystems. Positive impacts of precipitation on annual GEP have been reported for vegetated areas worldwide. However, little is known about the influence of precipitation intensity on GEP, especially at the monthly to seasonal temporal scale. Here we show that monthly GEP is insensitive to the sum of monthly total precipitation ( $P_s$ , mm), but positively correlated to precipitation intensity ( $P_a$ , mm), defined as the average precipitation per event from half-hourly measurements over a month. Different plant functional types (PFTs) exhibit substantial differences in the sensitivity of monthly GEP to  $P_a$ . PFTs of water-limited regions responded more intensely than those in mesic environments, as demonstrated by a negative correlation between the slope of the GEP- $P_a$  regression line and average  $P_a$ . Furthermore, this slope increases with latitude, indicating higher sensitivity of GEP to  $P_a$  for boreal ecosystems than for temperate regions. Therefore, we anticipate increased intensity of storms, as projected by some climate models, may impart a previously overlooked positive impact on precipitation intensity on GEP.

© 2012 Elsevier B.V. All rights reserved.

## 1. Introduction

Terrestrial ecosystems play a dynamic role in the global carbon (C) cycle as their carbon balance is highly sensitive to climatic change (Knapp and Smith, 2001; Piao et al., 2008). Climate variation affects photosynthesis and respiration processes in vegetation and therefore, climate is a major determinant of vegetation production of terrestrial ecosystems. As climatic and atmospheric changes are expected to accelerate in the near future, it is imperative that we improve the capability of estimating these factors (e.g., temperature, precipitation) on carbon budgets in support of scientific investigation and policy formulation (Chen et al., 2003).

While climatic change such as the increase in temperature and the elevated CO<sub>2</sub> concentration in the atmosphere, has been demonstrated to have large effects on terrestrial ecosystem production (Norby et al., 2005; Piao et al., 2008; Beer et al., 2010; Zhao and Running, 2010; Wu et al., 2012a), precipitation has been suggested to have an even profound impact on ecosystem dynamics than the single or combined effects of rising CO<sub>2</sub> and temperature, especially in arid and semiarid environments for the determination of the distribution, structure, and diversity of plants (Easterling et al., 2000; Houghton et al., 2001; Knapp et al., 2002; Weltzin et al., 2003).

The spatial pattern of precipitation in future, as projected by most general circulation models (GCMs), will increase in tropics and at middle and high latitudes and decrease in subtropical latitudes (Easterling et al., 2000). However, unresolved discrepancies exist among diverse models for specific regions, probably because of the difficulty in dealing with cloud microphysics and precipitation processes (Weltzin et al., 2003). Despite the uncertainties in feedbacks between annual precipitation variability and vegetation productivity, there are evidences that projected variation in future precipitation patterns are likely to alter the responses of the carbon cycle, though it may differ across biomes (Fang et al., 2001; Knapp and Smith, 2001; Wu and Chen, 2012; Wu et al., 2012b).

The underlying mechanism of the impacts of annual variability of precipitation on CO<sub>2</sub> uptake may differ across plant functional types (PFTs) and is difficult to evaluate globally due to the unavailability of data and model limitations. In addition, previous research conducted at the annual or seasonal temporal scales may have not fully addressed the effects of precipitation on vegetation biogeochemistry as precipitation is an environmental factor of high temporal heterogeneity (Chahine, 1992; Weltzin et al., 2003). Future precipitation regimes are characterized by larger sizes while also longer interval dry periods, it is thus important to understand how vegetation productivity responds to different frequencies and intensities of precipitation and how the sensitivities of such responses vary across different PFTs. Therefore, multi-year observations of study sites with long-term observations of productivity and precipitation are especially useful in

\* Corresponding author at: Department of Geography, University of Toronto, 100 St. George St., Room 5047, Toronto, ON, Canada M5S 3G3.

E-mail address: [hefery@163.com](mailto:hefery@163.com) (C. Wu).

quantifying the influence of long-term climate change on carbon sequestration of terrestrial ecosystems (Knapp and Smith, 2001; Huxman et al., 2004). In this study, we provide an analysis of the relationship between the gross ecosystem production (GEP) and monthly precipitation patterns using multi-year data (CO<sub>2</sub> flux and meteorological measurements) across eight PFTs in North America and ask:

- 1) How are monthly precipitation quantity and intensity related?
- 2) What is the relationship between monthly GEP and precipitation quantity and intensity?
- 3) How do these relationships vary by site characteristics?

## 2. Methods

### 2.1. Study sites

The data employed in this study were selected from the AmeriFlux (<http://public.ornl.gov/ameriflux/dataproducts.shtml>) and Fluxnet-Canada (<http://www.fluxnet-canada.ca>) archives. The rules for site selection were mainly regulated by the data availability, data quality and data time duration (e.g., minimum number of years of data was set to six) as these criteria can better support the mechanical analysis of long-term precipitation on productivity. As a result, data from 20 AmeriFlux and Fluxnet-Canada tower sites located across North America were chosen (Fig. 1 and Table 1). These sites cover eight PFTs in North America, including grassland (GRA, 2 sites), wetland (WET, 2 sites), deciduous broadleaf forest (DBF, 4 sites), crops (CRO, 2 sites), evergreen needle-leaf forest (ENF, 4 sites), mixed forest (MF, 2 sites), evergreen


broadleaf forest (EBF, 2 sites), and woody savannas (WSA, 2 sites). This broad range of sites provided a wide dynamical range of both GEP and precipitation. While there are a larger number of AmeriFlux sites available, the sites selected here have a long gap-filled time series of GEP and coincident high quality precipitation and volumetric soil moisture data collected on-site. Broadly, the selected sites are representative of many North American biomes.

### 2.2. Flux, climate and canopy structure data

For Canadian sites, a standard procedure was used to estimate annual net ecosystem production (NEP) and to partition NEP into components of GEP and ecosystem respiration (R<sub>e</sub>) from gap-filled half-hourly measurements (Barr et al., 2004). For AmeriFlux sites, level-4 GEP and products were used and these data were gap-filled with the Artificial Neural Network (ANN) method (Papale and Valentini, 2003) and/or the Marginal Distribution Sampling (MDS) method (Reichstein et al., 2005).

Previous work of Desai et al. (2008b) indicated that different decomposition techniques generally have a moderate impact on the modeled GEP, though most methods tended to cluster on similar results to within 10%, suggesting the reliability of multiple site comparisons and syntheses in this study when using a consistent method across all sites. This also agrees with the general understanding that biases associated with different gap filling methods tend to be small (Moffat et al., 2007).

Precipitation data for the sites were obtained from half-hourly meteorological measurements collected by on-site tipping bucket



**Fig. 1.** Locations of the 20 flux sites in North America that cover eight PFTs. GRA, WET, DBF, CRO, ENF, MF, EBF, and WSA, represent grassland, wetland, deciduous broadleaf forest, crops, evergreen needle-leaf forest, mixed forest, evergreen broadleaf forest, and woody savannas, respectively.

**Table 1**  
Detail descriptions of 20 sites in this analysis.

Sites	Latitude	Longitude	Land cover	Year	References
US-VAR	38.4133	−120.9507	GRA	2002–2007	Ryu et al. (2008)
CA-GRA	49.4300	−112.5600	GRA	1998–2006	Flanagan and Johnson (2005)
CA-FEN	54.9538	−112.4669	WET	2004–2009	Flanagan and Syed (2011)
CA-BOG	45.4094	−75.5187	WET	2004–2009	Admiral and Lafleur (2007)
US-UMB	45.5598	−84.7138	DBF	1999–2006	Curtis et al. (2002)
US-MMS <sup>a</sup>	39.3231	−86.4131	DBF	1999–2006	Dragoni et al. (2007)
US-WCR	45.8059	−90.0799	DBF	1999–2006	Cook et al. (2004)
CA-OA	53.6289	−106.1978	DBF	2002–2008	Barr et al. (2004)
US-NE3	41.1797	−96.4397	CRO	2002–2008	Suyker and Verma (2008)
US-BO1 <sup>b</sup>	40.0062	−88.2904	CRO	1997–2006	Meyers et al. (2006)
US-NR1	40.0329	−105.5460	ENF	2000–2007	Yi et al. (2008)
CA-OBS	53.9872	−105.1178	ENF	1999–2008	Margolis et al. (2006)
CA-OJP	53.9163	−104.6920	ENF	2000–2007	Margolis et al. (2006)
CA-QCJP	49.2671	−74.0365	ENF	2002–2009	Margolis et al. (2006)
US-SYV	46.2420	−89.3477	MF	2001–2006	Desai et al. (2008a)
US-HA1	42.5378	−72.1715	MF	2001–2006	Urbanski et al. (2007)
US-SP3	29.7548	−82.1633	EBF	1999–2004	Powell et al. (2008)
US-KS2	28.6086	−80.6715	EBF	2001–2006	Powell et al. (2008)
US-TON	38.4316	−120.9660	WSA	2002–2007	Ma et al. (2007)
US-SP2	29.7648	−82.2448	WSA	1999–2004	Gholz and Clark (2002)

<sup>a</sup> Missing data for 2004.

<sup>b</sup> Missing data for 2000 (GRA, WET, DBF, CRO, ENF, MF, EBF, and WSA, represent grassland, wetland, deciduous broadleaf forest, crops, evergreen needle-leaf forest, mixed forest, evergreen broadleaf forest, and woody savannas, respectively).

sensors. Missing data was gap-filled using the observations from nearby climate stations (Jassal et al., 2009). These data were summed to derive monthly precipitation ( $P_s$ , mm). We also calculated the number of time periods with precipitation ( $n$  = number of half-hours with recorded precipitation in the month) and the maximum precipitation of a single rain event ( $P_{max}$ , mm) from the half-hourly observations. These values were then used to obtain our estimate of precipitation intensity or average precipitation per rainfall event ( $P_a$ , mm) by dividing  $P_s$  by  $n$ .

Three further steps were adopted for preparing the monthly precipitation and GEP data. First, by the definition of  $P_a$ , months with no precipitation for all sites were excluded from the analysis. Second, for each site, months with differences between individual  $P_a$  and monthly average  $P_a$  larger than three times of the standard deviation (SD) of all  $P_a$  for this site were deemed as “outliers” (Ruan et al., 2005). Third, for all PFTs, we were focused on the growing season which was constrained by months with positive values in the mean air temperature ( $T_a$ ). However, due to differences in this growing season length as well as data availability across PFTs, certain criteria were also adopted. For example, for GRA and CRO, months from May to September were used (Flanagan and Johnson, 2005; Suyker and Verma, 2008), while months from April to October were excluded for Mediterranean sites, such as US-VAR and US-TON (Ma et al., 2007; Ryu et al., 2008).

For analysis of relationships between GEP and  $P_a$  or  $P_s$ , we also analyzed site-level soil moisture. Soil water content (SWC, %) was measured by the time-domain reflectometry (TDR) type probes at 0–30 cm for AmeriFlux sites. For Canadian sites, SWC was calculated as the average values of sampled data by depth to at least 50 cm. The variation of soil water content (VSWC) in  $i$  month is defined as the difference of mean SWC in current month and the previous month,

$$VSWC_i = SWC_i - SWC_{i-1}. \quad (1)$$

The difference of soil water content (DSWC) was calculated as

$$DSWC_i = \text{abs}(VSWC_i). \quad (2)$$

Finally, we further analyzed our relationships against maximum measured leaf area. We also searched in the above two data systems and literature sources for leaf area index (LAI,  $\text{m}^2\text{m}^{-2}$ ) data. At some sites, time series of monthly LAI was derived either as the average value from multiple observations (if available) or a single measurement per month.


### 3. Results

#### 3.1. Relationship between $P_s$ and $P_a$


Monthly observations of precipitation total ( $P_s$ ) and precipitation intensity ( $P_a$ ) were compared for all sites (Fig. 2). In general,  $P_a$  increases as  $P_s$  increases, with a linear fit between the two variables having a  $R^2$  of 0.24 ( $p < 0.001$ ). Some non-linearity was present in the end and the fit was improved with such a fit with coefficient of determination ( $R^2$ ) of 0.27 ( $p < 0.001$ ). The non-linear relationship indicates that  $P_a$  had a larger relative dynamical range than  $P_s$ . The importance of larger relative dynamic range for  $P_a$  is demonstrated in Fig. 3, which shows the averaged values of  $P_s$  and  $P_a$  for the eight PFTs that span wide ranges of monthly GEP and precipitation. For example, while no difference in  $P_s$  was observed between mixed forest (MF) and evergreen broadleaf forest (EBF) (91.61 and 91.67 mm, respectively),  $P_a$  varied considerably (0.89 mm and 1.61 mm, respectively), which correlates well with the large variation in GEP for these two groups (125 and 157  $\text{g C}/\text{m}^2/\text{month}$ , respectively). In this perspective, statistically speaking,  $P_a$  is likely to provide a better predictor of GEP than  $P_s$  and a thus greater potential for characterizing temporal and spatial patterns of vegetation production.

#### 3.2. Relationship between monthly GEP and $P_s$ , $P_a$

On an individual site basis, only three sites (CA-OBS, CA-OJP and US-SP3) had significant correlations between GEP and  $P_s$  (Fig. 4). This result indicates that this relationship differs from its counterpart at the annual scale where GEP is significantly associated with precipitation over 40% of the vegetated land, although with substantial variations across PFTs and spaces (Beer et al., 2010). In contrast, we find that monthly GEP was significantly correlated with  $P_a$  for all sites in our sample, with coefficients of determination ( $R^2$ ) ranging from 0.59 ( $p < 0.001$ ) to 0.09 ( $p = 0.012$ ) (Fig. 5).



**Fig. 2.** Relationship between  $P_s$  and  $P_a$  for the eight plant functional types. The regression line for overall data is  $y = 1.82 \exp [(2.24e^{-3})x] - 1.07$ . GRA, WET, DBF, CRO, ENF, MF, EBF, and WSA, represent grassland, wetland, deciduous broadleaf forest, crops, evergreen needle-leaf forest, mixed forest, evergreen broadleaf forest, and woody savannas, respectively.



**Fig. 3.** Average values of monthly GEP,  $P_s$  and  $P_a$  for the eight PFTs, GRA, WET, DBF, CRO, ENF, MF, EBF, and WSA, represent grassland, wetland, deciduous broadleaf forest, crops, evergreen needle-leaf forest, mixed forest, evergreen broadleaf forest, and woody savannas, respectively.

We substantiate this newly discovered relationship between GEP and  $P_a$  from both the canopy structure and the hydrologic perspectives. First, LAI is typically correlated to GEP, and here we find that similarly  $P_a$  is a better predictor of LAI variability than  $P_s$  (Fig. 6a, b). While a significant correlation ( $R^2$  equals to 0.26) is observed between  $P_a$  and LAI for all sites, the correlation differs across PFTs. The linear regressions are found to be the best fit for sites with relatively low LAI (e.g., GRA). However, for dense canopies (e.g., the EBF), the relationship breaks down at LAI greater than 6 (Table 2).

Second, the fate of precipitation includes interception, infiltration, and runoff (Chahine, 1992). In vegetated areas, a considerable part of the precipitation may be intercepted by the canopy and returned to the atmosphere via rapid evaporation. The remainder reaches the ground surface as throughfall and stemflow. Soil moisture is the direct link between precipitation and ecological processes, and only that part of precipitation that reaches the soil, i.e. net precipitation, can increase the soil moisture, either in the top layer or the entire root zone by hydrologic redistribution (Reynolds et al., 1999; Ryel et al., 2003; Gash and Shuttleworth, 2007; Muzyllo et al., 2009). For sites where SWC data were available ( $N=55\%$ ), we find that GEP is independent of mean SWC but negatively correlated to the monthly variation of the absolute difference in soil water content (DSWC, %) across PFTs,  $R^2=0.10$ ;  $p<0.001$  (Fig. 6c, d). Despite the small  $R^2$ , this finding indicates that vegetation growth likely benefits from a relatively stable SWC (e.g., a variation within 5%, Fig. 6f) at the monthly temporal scale because water typically reduces GEP during drought conditions, whereas when water is sufficient radiation availability is typically the limiting factor on GEP.

### 3.3. Sensitivity analysis of relationship between GEP and $P_a$

It is apparent from Fig. 5 that the sensitivity of GEP to  $P_a$  varies among the PFTs. Factors underlying this different sensitivity are explored in Fig. 7. The slopes derived from the linear regression for each biome were found to be negatively correlated with  $P_a$  ( $R^2=0.48$ ,  $p<0.001$ ), indicating that an increase in  $P_a$  has greater influence on GEP of PFTs in xeric regions than that in mesic environments (Fig. 7a). This result is consistent with the evolutionary and ecological views that species in water-limited regions with high production potentials (e.g., GRA, WET) would be very sensitive to the water availability, while biomes in mesic environments with high production potentials (e.g., EBF) are less sensitive (Huxman et al., 2004). Such different sensitivities are indirectly supported by recent research of both single site evaluations in mixed deciduous forest (Newman et al., 2006) and grasslands (Thomey et al., 2011) and multi-site studies across PFTs (Zha et al., 2010).

Dependence of the changes in this sensitivity on DSWC further supports these differences. Due to the lack of available SWC data

some sites were excluded from the analysis (CA-FEN, US-UMB, US-NE3, US-HA1, US-SP3 and US-SP2). Water limited PFTs, GRA for example, are characterized by the low  $P_a$  that leads to large variations in both DSWC and monthly GEP. However, for regions with high  $P_a$  (e.g., EBF), DSWC is small, resulting in a relatively stable monthly GEP. The positive correlation between DSWC and the coefficient of variation (CV) of GEP for sites with available SWC supports our analysis (Fig. 7b;  $R^2=0.55$ ,  $p<0.001$ ). This is because when water becomes abundant, resources other than water (e.g., soil nitrogen, light) may act as more important limiting factors for production (Huxman et al., 2004). These two different responses imply a decrease in the amplitude of GEP variation at high  $P_a$  and explain the negative correlation between the slope and  $P_a$ .

We further analyzed GEP- $P_a$  sensitivity as a function of latitudinal gradient. A coefficient of determination ( $R^2$ ) of 0.49 ( $p<0.001$ ) was obtained between the slopes of the GEP- $P_a$  relationship and latitude for all sites (Fig. 8a), implying boreal ecosystems have a higher sensitivity than that of temperate regions.  $P_a$  is also better correlated with GEP of sites located at high latitudes, which is evidenced by the relationship between  $R^2$  of GEP- $P_a$  relationship and latitude (Fig. 8b). A negative correlation between  $R^2$  and average monthly  $P_s$  for all sites (Fig. 8c;  $R^2=0.57$ ,  $p<0.001$ ) highlights that the use of  $P_a$  to evaluate future precipitation intensity on GEP would be more suitable for drought-prone regions.

### 3.4. Precipitation intensity covariation with air temperature and global shortwave radiation

Since temperature and radiation are important factors of GEP, we also analyzed their covariance with precipitation intensity (Table 3). We found that monthly mean temperature was significantly correlated with  $P_a$  except for the grassland sites and the coefficients of determination also showed substantial variations among PFTs. WET ecosystem had the highest correlation with  $R^2$  of 0.47 ( $p<0.001$ ) and the lowest  $R^2$  of 0.10 ( $p=0.025$ ) was observed at CRO sites. Relationship between  $P_a$  and  $R_g$  was much lower (below 0.20) compared to that of  $P_a$  and  $T_a$  and they were only significantly correlated at WET, MF, EBF and WSA ecosystems. These analyses indicate that for most of these ecosystems, the temperature may co-control the vegetation production with the precipitation intensity. However, the influence of radiation is much smaller than that of temperature.

## 4. Discussion

To the best of our knowledge, these are the first results showing the additional important and overlooked impact of precipitation intensity on GEP at the monthly temporal scale. These results are consistent with previous research showing that while most aspects of terrestrial ecosystem structure and function are vulnerable to hydrologic changes caused by precipitation, the response may be less dependent on changes in annual precipitation quantity but more reliant on precipitation variability (Mearns et al., 1997; Knapp et al., 2002).  $P_a$  in our analysis, although being derived from the total half-hourly measurements, holds the potential as an indicator of the precipitation variability that best relates to GEP variability. These results are also consistent with previous experimental work altering rain amount in grasslands (Heisler-White et al., 2008) and observational studies (Thomey et al., 2011), which report a significant increase of above ground net primary production (ANPP) for grasslands with a single large rainfall event compared to multiple small events with equal total rainfall amounts.

Our analyses suggest that there is a relationship between maximum production and VSWC. Although not fully understood yet, the function between monthly GEP and VSWC can be conceptually illustrated by Fig. 9. Soil water content stability means values may fall in the  $[-5, 5]$  for example, so why do these samples have a relatively higher monthly GEP? Large negative values (left region of VSWC =  $-5\%$ ) mean




Fig. 4. Relationship between the monthly sum of precipitation ( $P_s$ , mm) and monthly GEP ( $g \cdot C \cdot m^{-2}$ ) for 20 sites in North America. a, b, c, d, e, f, g and h represent the GRA, WET, DBF, CRO, ENF, MF, EBF, and WSA, respectively (see the text for details).





Fig. 5. Relationship between the monthly average precipitation per event ( $P_a$ , mm) and monthly GEP ( $\text{g} \cdot \text{C} \cdot \text{m}^{-2}$ ) for 20 sites in North America. a, b, c, d, e, f, g and h represent the GRA, WET, DBF, CRO, ENF, MF, EBF and WSA, respectively (see the text for details).



**Fig. 6.** Relationships between precipitation patterns and monthly GEP, canopy structural and soil water variables. a, correlation between leaf area index (LAI) and  $P_s$ . b, correlation between leaf area index (LAI) and  $P_a$ . c, correlation between monthly GEP and the mean soil water content (SWC). d, correlation between monthly GEP and the DSWC. e, correlation between  $P_s$  and the DSWC. f, correlation between  $P_a$  and the DSWC. GRA, WET, DBF, CRO, ENF, MF, EBF, and WSA, represent grassland, wetland, deciduous broadleaf forest, crops, evergreen needle-leaf forest, mixed forest, evergreen broadleaf forest, and woody savannas, respectively.

drought and the decrease of monthly GEP is caused by the water shortage. On the other hand, large positive variations (right region of  $VSWC=5\%$ ) in SWC are caused by the increase in the number of periods of precipitation ( $n$ ). However, increase in  $n$  will also result to low radiation which can also reduce the monthly GEP. These assumptions are supported by our data (Fig. 10). Both low and high values of  $n$  can lead to large VSWC.

Average  $n$  ( $n_{ave}$ ) for VSWC below  $-5\%$  and above  $5\%$  are 56 and 102, respectively. Months with VSWC variations within  $5\%$  have a medium  $n_{ave}$  of 70. However, high  $n$  values can give rise to sharp decreases in the radiation for all observations with  $R^2$  equals to 0.20 ( $p<0.001$ ). Previous research also suggests that water availability will play a limiting role for production in drought conditions, while other resources


**Table 2**  
Correlations between  $P_a$  and available LAI across plant functional types.

PFTs	Equations	R <sup>2</sup> (p-value)	No. (n)	LAI_ave (m <sup>2</sup> ·m <sup>-2</sup> )	SD (m <sup>2</sup> ·m <sup>-2</sup> )
GRA	$L = 1.67P_a - 0.31$	0.42 (p<0.001)	37	0.87	0.65
DBF	$L = 1.68\ln P_a + 3.87$	0.26 (p<0.001)	43	3.72	1.18
CRO	$L = 1.64\ln P_a + 1.62$	0.56 (p<0.001)	26	2.11	1.26
ENF	$L = 0.21\ln P_a + 2.16$	0.08 (p=0.781)	12	2.25	0.65
MF	$L = 4.08P_a - 0.53$	0.58 (p=0.006)	11	2.83	1.68
EBF	$L = 0.85\ln P_a + 4.90$	0.26 (p<0.001)	63	5.14	0.84
WSA	$L = 1.10P_a + 0.57$	0.41 (p<0.001)	63	2.17	1.78


(SD is the standard deviation; GRA, DBF, CRO, ENF, MF, EBF, and WSA, represent grassland, deciduous broadleaf forest, crops, evergreen needle-leaf forest, mixed forest, evergreen broadleaf forest, and woody savannas, respectively).

(e.g., light, nitrogen) will become the constraint role when water is sufficient (Weltzin et al., 2003). Therefore, samples with a medium VSWC are expected to enhance monthly GEP.

This conceptual model is also useful in explaining the positive correlation between  $P_a$  and GEP. In natural environments, it is difficult to maintain the desired stable state of VSWC because of constant changes in environmental factors (precipitation, evaporation, etc.). A significant logarithmic decrease of  $P_a$  with DSWC is found from the available data (as shown in previous Fig. 6f), suggesting that larger  $P_a$  values promote a more stable net precipitation, which leads to a relatively more stable SWC, supporting enhanced GEP. This reasoning is



**Fig. 7.** a, Correlation between  $P_a$  and the slope of GEP- $P_a$  relationship for all sites; b, Correlation between DSWC and the coefficient of variation (CV) of GEP for sites with available SWC (CA-FEN, US-UMB, US-NE3, US-HA1, US-SP3 and US-SP2 are excluded due to unavailable of SWC data).



**Fig. 8.** Latitudinal patterns of (a) the slope of GEP- $P_a$  correlation and (b)  $R^2$  of GEP- $P_a$  correlation for all sites; (c) relationship between  $R^2$  of GEP- $P_a$  correlation and average  $P_s$  for all sites.

supported by our available datasets showing that less than 6% (5 out of 88) observations have SWC variations over 5% for  $P_a$  above 1.5 mm, whereas for  $P_a$  below 1 mm this proportion rises to 31% (68 out of 221). We believe that it is this result that best explains the observed positive influence of increased precipitation intensity on monthly GEP.

Since  $P_a$  is related to both LAI and GEP, it is also worthwhile to consider whether variation in LAI over the growing season would potentially affect the correlation between  $P_a$  and GEP. We standardized GEP by LAI using the ratio GEP/LAI as one way to remove this effect. Detailed monthly LAI data were available at five sites, and the correlations



**Table 3**

Relationships between precipitation intensity ( $P_a$ , mm) and monthly mean air temperature ( $T_a$ , °C), global shortwave radiation ( $R_g$ , MJ/m<sup>2</sup>/month).

Plant functional types		$T_a$ (°C)	$R_g$ (MJ/m <sup>2</sup> /month)
$P_a$ (mm)	GRA	NS	NS
	WET	$R^2=0.47$ , $p<0.001$	$R^2=0.11$ , $p=0.005$
	DBF	$R^2=0.27$ , $p<0.001$	NS
	CRO	$R^2=0.10$ , $p=0.025$	NS
	ENF	$R^2=0.36$ , $p<0.001$	NS
	MF	$R^2=0.35$ , $p<0.001$	$R^2=0.18$ , $p<0.001$
	EBF	$R^2=0.35$ , $p<0.001$	$R^2=0.12$ , $p<0.001$
	WSA	$R^2=0.34$ , $p<0.001$	$R^2=0.16$ , $p<0.001$

GRA, WET, DBF, CRO, ENF, MF, EBF, and WSA represent grassland, wetland, deciduous broadleaf forest, crops, evergreen needle-leaf forest, mixed forest, evergreen broadleaf forest, and woody savannas, respectively. NS indicates no significant correlation was observed.

between  $P_a$  and GEP/LAI for these sites are shown in Fig. 11. While positive correlation was obtained at the grassland site (CA-GRA,  $R^2=0.11$ ,  $p<0.001$ ) and the deciduous forest site (US-MMS,  $R^2=0.21$ ,  $p<0.001$ ), both the crop site (US-NE3) and woody savannas site (US-SP2) have shown negative correlations with both  $R^2$  of 0.31 ( $p<0.001$ ). No correlation was obtained for an evergreen broadleaf forest site (US-SP3). The underlying mechanism of these equivocal correlations is not well understood at present, but still suggests the influence of precipitation intensity on vegetation productivity and should be explored in future analysis.

Our evaluation on the relationships between  $P_a$  and temperature and radiation shows that the effect of  $P_a$  on monthly GEP should be considered with other factors influencing GEP. This agrees with the assumption that the roles of climate variables on vegetation production could change with both time and PFTs. More importantly, it supports the suggestion that the pattern of precipitation (e.g., intensity) could have more important effect than the quantity. This is meaningful for the regional carbon budget since we have many cases where predictions of vegetation production are substantially different among various ecosystem models (Beer et al., 2010). One explanation of such difference could be the unknown effects of certain variables (e.g., precipitation patterns) that affect GEP, but their mechanisms are not well understood and thus not being included in models. Our results from precipitation intensity may be a potential solution to know the missing processes or feedback mechanisms that attenuate the vegetation production to climate (Beer et al., 2010). Given the difficulty in climate models predicting future precipitation, one particular useful implication of our analysis is that future ecosystem models for the analysis of precipitation effects might take this precipitation intensity into consideration to improve the ability of climate models to predict quantitative ecosystem responses in future climate scenarios. However, for the operational




Fig. 9. The conceptual model of maximum production as a function of VSWC.




Fig. 10. a, Relationship between precipitation times ( $n$ ) and the variation in soil water content (VSWC). The two gray lines indicate the variations with  $\pm 5\%$ . Black line represents the regression equation. Average  $n$  ( $n_{ave}$ ) for VSWC below  $-5\%$ , within  $5\%$  and above  $5\%$  are 56, 70 and 102, respectively. b, Relationship between the precipitation times ( $n$ ) and the monthly global shortwave radiation.

applications of  $P_a$ , the effects of other meteorological variables (e.g., temperature, radiation) on vegetation production should be also considered as their covariation with precipitation intensity.

Two extreme conditions also need to be discussed in the application of  $P_a$ . First, for months with no precipitation,  $P_a$  will not make sense for the evaluation of water availability. The production of vegetation can either be very low (US-VAR) or even higher (US-TON) and it may be dependent on the synchronization between vegetation growing season and precipitation period. In this case, other factors, such as the soil water content, soil texture and the precipitation of the last month may have potential use for the analysis of water status and production (Ryel et al., 2003; Weltzin et al., 2003). A second consideration is the “outlier” months that have  $P_a$  departing far from the average values that are often due to unusual and rarely occurring storms. For example,  $P_a$  for site US-BO1 reached 9.5 mm in September 2004 because there were only two ( $n=2$ ) rainfalls that occurred in that month and one of which had a precipitation amount of 18 mm ( $P_{max}=18$  mm), almost equal to  $P_s$  (19 mm). This unusual  $P_a$  value is almost five times larger than the SD of  $P_a$  after subtracting the  $P_a$  averages of all months for this site. This “outlier”, in a statistical perspective, would happen in natural ecosystems because precipitation is a stochastic environmental factor that can change dramatically. However, for the month, both GEP and soil water content are not




Fig. 11. Relationship between  $P_a$  and GEP/LAI for (a) CA-GRA, (b) US-MMS, (c) US-NE3, (d) US-SP3 and (e) US-SP2.

significantly different from other months of the same period because high values of  $P_a$  may not imply large amount of monthly precipitation. This uncertainty points to the need of incorporation of other variables (e.g., precipitation intervals) in order to better characterize the frequency and intensity of precipitation events.

### 5. Conclusions

There are several broad implications of this newly developed relationship between our precipitation intensity metric ( $P_a$ ) and gross ecosystem photosynthesis (GEP). First,  $P_a$  highlights the importance of precipitation temporal patterns rather than the quantity alone. It provides insights to improve the accuracy of GEP estimates by ecosystems models that do not incorporate such mechanisms. Second, the correlation between GEP and  $P_a$  can bridge previous experimental results on carbon components under manipulated precipitation patterns to natural settings, in order to improve our understanding of the role of precipitation variability (Knapp et al., 2002). Third, this

correlation provides a method to link GEP to changes of precipitation at the monthly temporal scale. Previous results that are obtained at the annual scale may be improved using  $P_a$ , which would be especially useful in climate change research because temporal heterogeneity of precipitation can lead to substantial differences in monthly precipitation even if the annual precipitation totals are relatively stable. Lastly, the higher correlation coefficients at water limited sites indicate that  $P_a$  would be most useful for the arid, semi-arid and drought susceptible regions that occupy about 30% of the world's land surface and support about one billion people (Weltzin et al., 2003). As projected by climate change models, drought is likely to become more severe at subtropical latitudes, thus we may expect that the area suitable for  $P_a$  application will increase. There is an urgent need for models to assess this dependence of vegetation production on precipitation patterns at different temporal scales and substantiate our claim of superior performance of  $P_a$  over  $P_s$  under relatively dry climate conditions and its importance in estimating the global terrestrial productivity and carbon cycle under future climate change scenarios.

## Acknowledgments

This work used the data at the sites from both the AmeriFlux and Fluxnet-Canada. We are grateful for the PIs of these sites in providing the data and explanations and acknowledging the funding sources, including US DOE, NSF, and Environment Canada that supported these observations. This work was funded by an NSERC Strategic Grant (381474–09), the National Natural Science Foundation (grant no. 41001210), and the Knowledge Innovation Program of CAS (KZCX2-EW-QN302).

## References

- Admiral, S.W., Lafleur, P.M., 2007. Modelling of latent heat partitioning at a bog peatland. *Agricultural and Forest Meteorology* 144, 213–229.
- Barr, A.G., Black, T.A., Hogg, E.H., Kljun, N., Morgenstern, K., Nesic, Z., 2004. Interannual variability in the leaf area index of a boreal aspen-hazelnut forest in relation to net ecosystem production. *Agricultural and Forest Meteorology* 126, 237–255.
- Beer, C., Reichstein, M., Tomelleri, E., Ciais, P., Jung, M., Carvalhais, N., Rödenbeck, C., Arain, M.A., Baldocchi, D., Bonan, G.B., Bondeau, A., Cescatti, A., Lasslop, G., Lindroth, A., Lomas, M., Luysaert, S., Margolis, H., Oleson, K.W., Rouspard, O., Veenendaal, E., Viovy, N., Williams, C., Ian Woodward, F., Papale, D., 2010. Terrestrial gross carbon dioxide uptake: global distribution and covariation with climate. *Science* 329, 834–838.
- Chahine, M.T., 1992. The hydrological cycle and its influence on climate. *Nature* 359, 373–380.
- Chen, J.M., Ju, W., Cihlar, J., Price, D., Liu, J., Chen, W., Pan, J., Black, T.A., Barr, A., 2003. Spatial distribution of carbon sources and sinks in Canada's forests based on remote sensing. *Tellus B* 55, 622–642.
- Cook, B.D., Davis, K.J., Wang, W., Desai, A., Berger, B.W., Teclaw, R.M., Martin, J.G., Bolstad, P.V., Bakwin, P.S., Yi, C., Heilman, W., 2004. Carbon exchange and venting anomalies in an upland deciduous forest in northern Wisconsin, USA. *Agricultural and Forest Meteorology* 126, 271–295.
- Curtis, P.S., Hanson, P.J., Bolstad, P., Barford, C., Randolph, J.C., Schmid, H.P., Wilson, K.B., 2002. Biometric and eddy-covariance based estimates of annual carbon storage in five eastern North American deciduous forests. *Agricultural and Forest Meteorology* 113, 3–19.
- Desai, A.R., Noormets, A., Bolstad, P.V., Chen, J.Q., Cook, B.D., Davis, K.J., Euskirchen, E.S., Gough, C.M., Martin, J.G., Ricciuto, D.M., Schmid, H.P., Tang, J.W., Wang, W.G., 2008a. Influence of vegetation and seasonal forcing on carbon dioxide fluxes across the Upper Midwest, USA: implications for regional scaling. *Agricultural and Forest Meteorology* 148, 288–308.
- Desai, A.R., Richardson, A.D., Moffat, A.M., Kattge, J., Hollinger, D.Y., Barr, A., Falge, E., Noormets, A., Papale, D., Reichstein, M., Stauch, V.J., 2008b. Cross site evaluation of eddy covariance GPP and RE decomposition techniques. *Agricultural and Forest Meteorology* 148, 821–838.
- Dragoni, D., Schmid, H.P., Grimmond, C.S.B., Loescher, H.W., 2007. Uncertainty of annual net ecosystem productivity estimated using eddy covariance flux measurements. *Journal of Geophysical Research* 112, D17102 <http://dx.doi.org/10.1029/2006JD008149>.
- Easterling, D.R., Meehl, G.A., Parmesan, C., Changnon, S.A., Karl, T.R., Mearns, L.O., 2000. Climate extremes: observations, modeling, and impacts. *Science* 289, 2068–2074.
- Fang, J., Piao, S., Tang, Z., Peng, C., Ji, W., 2001. Interannual variability in net primary production and precipitation. *Science* 293, 1723.
- Flanagan, L.B., Johnson, B.G., 2005. Interacting effects of temperature, soil moisture and plant biomass production on ecosystem respiration in a northern temperate grassland. *Agricultural and Forest Meteorology* 130, 237–253.
- Flanagan, L.B., Syed, K.H., 2011. Stimulation of both photosynthesis and respiration in response to warmer and drier conditions in a boreal peatland ecosystem. *Global Change Biology* 17, 2271–2287.
- Gash, J.H.C., Shuttleworth, W.J., 2007. *Benchmark Papers in Hydrology: Evaporation*. IAHS Press, Wallingford, p. 521.
- Gholz, H.L., Clark, K.L., 2002. Energy exchange across a chronosequence of slash pine forests in Florida. *Agricultural and Forest Meteorology* 112, 87–102.
- Heisler-White, J.L., Knapp, A.K., Kelly, E.F., 2008. Increasing precipitation event size increases aboveground net primary productivity in a semi-arid grassland. *Oecologia* 158, 129–140.
- Houghton, J.T., Ding, Y., Griggs, D.J., Nogueira, M., van der Linden, P.J., Dai, X., Maskell, K., Johnson, C.A., 2001. *Climate Change 2001: The Scientific Basis Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge (United Kingdom).
- Huxman, T.E., Smith, M.D., Fay, P.A., Knapp, A.K., Shaw, M.R., Loik, M.E., Smith, S.D., Tissue, D.T., Zak, J.C., Weltzin, J.F., Pockman, W.T., Sala, O.E., Haddad, B., Harte, J., Koch, G.W., Schwinning, S., Small, E.A., Williams, D.G., 2004. Convergence across biomes to a common rain-use efficiency. *Nature* 429, 651–654.
- Jassal, R.S., Black, T.A., Spittlehouse, D.L., Brümmer, C., Nesic, Z., 2009. Evapotranspiration and water use efficiency in different-aged Pacific Northwest Douglas-fir stands. *Agricultural and Forest Meteorology* 149, 1168–1178.
- Knapp, A.K., Smith, M.D., 2001. Variation among biomes in temporal dynamics of aboveground primary production. *Science* 291, 481–484.
- Knapp, A.K., Fay, P.A., Blair, J.M., Collins, S.L., Smith, M.D., Carlisle, J.D., Harper, C.W., Danner, B.T., Lett, M.S., McCarron, J.K., 2002. Rainfall variability, carbon cycling and plant species diversity in a mesic grassland. *Science* 298, 2202–2205.
- Ma, S.Y., Baldocchi, D.D., Xu, L.K., Hehn, T., 2007. Inter-annual variability in carbon dioxide exchange of an oak/grass savanna and open grassland in California. *Agricultural and Forest Meteorology* 147, 157–171.
- Margolis, H.A., Flanagan, L.B., Amiro, B.D., 2006. The Fluxnet-Canada Research Network: influence of climate and disturbance on carbon cycling in forests and peatlands. *Agricultural and Forest Meteorology* 140, 1–5.
- Mearns, L.O., Rosenzweig, C., Goldberg, R., 1997. Mean and variance change in climate scenarios: methods, agricultural applications, and measures of uncertainty. *Climatic Change* 35, 367–396.
- Meyers, T.P., Luke, W.T., Meisinger, J.J., 2006. Fluxes of ammonia and sulfate over maize using relaxed eddy accumulation. *Agricultural and Forest Meteorology* 136, 203–213.
- Moffat, A.M., Papale, D., Reichstein, M., Richardson, A.D., Barr, A.G., Beckstein, C., Braswell, B.H., Churkina, G., Desai, A.R., Falge, E., Gove, J.H., Heimann, M., Hui, D., Jarvis, A.J., Kattge, J., Noormets, A., Stauch, V.J., 2007. Comprehensive comparison of gap-filling techniques for eddy covariance net carbon fluxes. *Agricultural and Forest Meteorology* 147, 209–232.
- Muzylo, A., Llorens, P., Valente, F., Keizer, J.J., Domingo, F., Gash, J.H.C., 2009. A review of rainfall interception modeling. *Journal of Hydrology* 370, 191–206.
- Newman, G.S., Arthur, M.A., Muller, R.N., 2006. Above- and belowground net primary production in a temperate mixed deciduous forest. *Ecosystems* 9, 317–329.
- Norby, R.J., DeLucia, E.H., Gielen, B., Calfapietra, C., Giardina, C.P., King, S.J., Ledford, J., McCarthy, B.H., Moore, D.J.P., Ceulemans, R., De Angelis, P., Finzi, A.C., Karnosky, D.F., Kubiske, M.E., Lukac, M., Pregitzer, K.S., Scarasci-Mugnozza, G.E., Schlesinger, W.H., Oren, R., 2005. Forest response to elevated CO<sub>2</sub> is conserved across a broad range of productivity. *Proceedings of the National Academy of Sciences* 102, 18052–18056.
- Papale, D., Valentini, A., 2003. A new assessment of European forests carbon exchange by eddy fluxes and artificial neural network spatialization. *Global Change Biology* 9, 525–535.
- Piao, S., Ciais, P., Friedlingstein, P., Peylin, P., Reichstein, M., Luysaert, S., Margolis, H., Fang, J., Barr, A., Chen, A., Grelle, A., Hollinger, D.Y., Laurila, T., Lindroth, A., Richardson, A.D., Vesala, T., 2008. Net carbon dioxide losses of northern ecosystems in response to autumn warming. *Nature* 451, 49–52.
- Powell, T.L., Gholz, H.L., Clark, K.L., Starr, G., Cropper, W.P., Martin, T.A., 2008. Carbon exchange of a mature, naturally regenerated pine forest in north Florida. *Global Change Biology* 14, 2523–2538.
- Reichstein, M., Falge, E., Baldocchi, D., Papale, D., Aubinet, M., Berbigier, P., Bernhofer, C., Buchmann, N., Gilmanov, T., Granier, A., Grunwald, T., Havrankova, K., Ilvesniemi, H., Janous, D., Knohl, A., Laurila, T., Lohila, A., Loustau, D., Matteucci, G., Meyers, T., Miglietta, F., Ourcival, J.M., Pumpanen, J., Rambal, S., Rotenberg, E., Sanz, M., Tenhunen, J., Seufert, G., Vaccari, F., Vesala, T., Yakir, D., Valentini, R., 2005. On the separation of net ecosystem exchange into assimilation and ecosystem respiration: review and improved algorithm. *Global Change Biology* 11, 1424–1439.
- Reynolds, J.E., Virginia, R.A., Kemp, P.R., Soya, A.G., Tremmel, D.C., 1999. Impact of drought on desert shrubs: effects of seasonality and degree of resource island development. *Ecological Monographs* 69, 69–106.
- Ruan, D., Chen, G., Kerre, E.E., Wets, G., 2005. *Intelligent Data Mining: Techniques and Applications*. Studies in Computational Intelligence, vol. 5. Springer, p. 318.
- Ryel, R.J., Caldwell, M.M., Leffler, A.J., Yoder, C.K., 2003. Rapid soil moisture recharge to depth by roots in a stand of *Artemisia tridentata*. *Ecology* 84, 757–764.
- Ryu, Y., Baldocchi, D.D., Ma, S., Hehn, T., 2008. Interannual variability of evapotranspiration and energy exchange over an annual grassland in California. *Journal of Geophysical Research* 113, D09104 <http://dx.doi.org/10.1029/2007JD009263>.
- Suyker, A.E., Verma, S.B., 2008. Interannual water vapor and energy exchange in an irrigated maize-based agroecosystem. *Agricultural and Forest Meteorology* 148, 417–427.
- Thomey, M.L., Collins, S.L., Vargas, R., Johnson, J.E., Brown, R.F., Natvig, D.O., Friggens, M.T., 2011. Effect of precipitation variability on net primary production and soil respiration in a Chihuahuan Desert grassland. *Global Change Biology* 17, 1505–1515.
- Urbanski, S., Barford, C., Wofsy, S., Kucharik, C., Pyle, E., Budney, J., McKain, K., Fitzjarrald, D., Czirkowsky, M., Munger, J.W., 2007. Factors controlling CO<sub>2</sub> exchange on timescales from hourly to decadal at Harvard Forest. *Journal of Geophysical Research* 112, G02020 <http://dx.doi.org/10.1029/2006JG000293>.
- Weltzin, J.F., Loik, M.E., Schwinning, S., Williams, D.G., Fay, P., Haddad, B., Harte, J., Huxman, T.E., Knapp, A.K., Lin, G., Pockman, W.T., Shaw, M.R., Small, E.E., Smith, M.D., Smith, S.D., Tissue, D.T., Zak, J.C., 2003. Assessing the response of terrestrial ecosystems to potential changes in precipitation. *BioScience* 53, 941–952.
- Wu, C., Chen, J.M., 2012. The use of precipitation intensity in estimating gross primary production in four northern grasslands. *Journal of Arid Environments* 82, 11–18.
- Wu, C., Chen, J.M., Gonsamo, A., Price, D.T., Black, T.A., Kurz, W.A., 2012a. Interannual variability of carbon sequestration is determined by the lag between ends of net uptake and photosynthesis: evidence from long records of two contrasting forest stands. *Agricultural and Forest Meteorology* 164, 29–38.
- Wu, C., Chen, J.M., Pumpanen, J., Cescatti, A., Marcolla, B., Blanken, P.D., Ardo, J., Tang, Y., Magliulo, V., Georgiadis, T., Soegaard, H., Cook, D.R., Harding, R.J., 2012b. An underestimated role of precipitation frequency in regulating summer soil moisture. *Environmental Research Letters* 7, 024011 <http://dx.doi.org/10.1088/1748-9326/7/2/024011>.
- Yi, C., Anderson, D.E., Turnipseed, A.A., Burns, S.P., Sparks, J., Stannard, D., Monson, K.R., 2008. The contribution of advective fluxes to net ecosystem CO<sub>2</sub> exchange in a high-elevation, subalpine forest ecosystems. *Ecological Applications* 18, 1379–1390.
- Zha, T., Barr, A.G., Kamp, G., Black, T.A., McCaughey, J.H., Flanagan, L.B., 2010. Interannual variation of evapotranspiration from forest and grassland ecosystems in western Canada in relation to drought. *Agricultural and Forest Meteorology* 150, 1476–1484.
- Zhao, M., Running, S.W., 2010. Drought-induced reduction in global terrestrial net primary production from 2000 through 2009. *Science* 329, 940–943.